


The Structure of the Earth	
The Crust	Varies in thickness (5-10km beneath the ocean). Amounts to less than 1% of the Earth's total mass. Made up of serval major plates.
The Mantle	Widest layer (2900km thick). The heat and pressure means the rock is in a liquid state that is in a state of convection.
The Inner and outer Core	Hottest section (5000 degrees). Mostly made of iron and nickel and is 4x denser than the crust. Inner section is solid whereas outer layer is liquid.



What is a Tectonic Plate?

A tectonic plate is a **massive, irregularly shaped slab of solid rock**, composed of both **continental** and **oceanic** lithospheres. These tectonic plates move in various ways against each other on areas know as plate margins.

Theory of Plate Tectonics

In 1912, Alfred Wegener proposed the theory of continental drift. He suggested the existence of Pangaea and that continents drift. Evidence for this includes;

1	Palaeomagnetism – Record of the Earth's polarity on erupted lava.
2	Geology - Rock sequences and jigsaw fitting of the world's continents.
3	Fossil records –Fossil remains of reptiles found in different continents.
4	Living species – Some species found on different continents are similar.
5	Climatology - Glacial deposits on the Equator suggests plate movement.



Types of Plate Boundaries

Divergent/Destructive Plate Boundaries

Oceanic – Continental: Subduction of an ocean plate at oceanic and continental plate margins leads to fold mountains & volcanoes.

Andean Mountain Range, Peru and Chile

Oceanic – Oceanic: When two oceanic plates collide the older and denser plate subducts. The process here creates volcanic island arcs such as those found in the Lesser Antilles.

Aleutian Island, Alaska USA

Continental - Continental: Involves two plate margins that are both continental and neither subducts. As these two plates are similar in density, the two plates collide to uplift and fold the crust.

Himalayan Mountain Range, Nepal and China

Divergent/Constructive Plate Boundaries

Continental – Continental: Caused by geologically recent mantle plume splitting a continental plate to create a new ocean basin. It can cause Basaltic volcanoes and minor earthquakes.

African Rift Valley, Ethiopia

Oceanic – Oceanic: New lithosphere forms at constructive margins, where rising plumes of magma stretches the crust to create intense volcanic activity on the ocean floor.

Mid-Atlantic Ridge, Atlantic Ocean

Conservative Plate Boundary

Oceanic – Continent: Two plates slide past each other in either different directions or the same direction but at different speeds. As they shear past they can cause powerful earthquakes.

San Andreas Fault, California USA

Volcanic Hotspots

A concentration of radioactive elements inside the mantle may cause a hotspot to develop. From this, a plume of magma rises to melt through into the plate above. Where lava breaks through to the surface, active volcanoes can occur above the hot spot.

Intra-plate Earthquake

An intra-plate earthquake refers to an earthquake that occurs within the interior of a tectonic plate.

Global Distribution of Tectonic Hazards

Earthquakes

Earthquakes occur throughout the world but predominately on **plate boundaries**. For example the San Andreas Fault, a conservative plate margin. Furthermore, earthquakes also occur on the constructive plate boundaries of the Mid- Atlantic Ridge, although these are not as severe when compared to conservative, collision and especially destructive plate margins.

Volcanoes

Volcanoes are most likely to occur along **subduction zones** where oceanic plates dive under continental plates. Volcanic activity can also be found along **constructive plate margins** such as the Mid Atlantic ridge. There are, however, exceptions. The Hawaiian Islands, which are entirely volcanic in origin, formed in the middle of the Pacific Ocean. This is explained by the **'hotspot' theory**.

Tsunamis

The global distribution of tsunamis is fairly predictable, with around 90% of all events occurring **within the Pacific Basin**, associated with activity at **plate margins**. Most are generated at **subduction zones**, particularly off the Japan-Taiwan island arc, South America and the Aleutian Islands.

What is the Asthenosphere?

The upper layer of the earth's mantle, below the lithosphere, in which there is relatively low resistance to plastic flow and convection is thought to occur.

Mechanism of Plate Movement

The lithosphere is divided into tectonic plates. The processes that cause their movement are still debated. Below are some of the up-to-date theories surrounding reasons why plates move.

Slab Pull	Newly formed oceanic lithosphere at mid ocean ridges is less dense than the asthenosphere, but becomes denser with age as it cools and thickens. This causes it to sink into the mantle at subduction zones (Mariana Trench), pulling slabs of lithosphere apart at divergent boundaries and resulting in sea floor spreading or rifting. This process linked to driving convection currents within the mantle.
Ridge Push	As the lithosphere formed at divergent plate margins is hot, and less dense that the surrounding area, it rises to form oceanic ridges (Mid Atlantic Ridge). The newly-formed plates slide sideways off these high areas, pushing the plate in front of them resulting in a ridge-push mechanism.

Dynamic Landscapes: Tectonic Processes & Hazards

Types of Lithospheric Plates

Continental	Oceanic
- Thick (10-70km) - Buoyant (less dense than oceanic crust) - Old sedimentary & metamorphic rock	- Thin (-7 km) - Dense (sinks under continental crust) - Young basalt (igneous) rock

Benioff Zone and Subduction Processes


The **Benioff Zone** is an inclined zone in which many deep earthquakes occur, situated beneath a destructive plate boundary where oceanic crust is being subducted.

As the **asthenosphere** and **lithosphere** at the ridge are heated, they expand and become elevated above the surrounding sea floor.

At a **subduction boundary**, one plate is denser and heavier than the other plate. The denser, heavier plate begins to **subduct** beneath the plate that is less dense.


The subducting plate is **much colder and heavier** than the mantle, so it continues to sink, pulling the rest of the plate along with it. The force that the sinking edge of the plate exerts on the rest of the plate is called **slab pull**.

Benioff Zone and Earthquakes



How do Earthquakes happen?

Earthquakes (shallow focus – less than 70km) happen when two plates become **locked** causing **friction** to build up. From this **stress**, the **pressure** will eventually be released, triggering the plates to move into a new position. This movement causes energy in the form of **seismic waves**, to travel from the **focus** towards the **epicentre**. As a result, the crust vibrates triggering an earthquake.



Types of Seismic Waves

P Waves	Travel through solids and liquids. Shakes the Earth in the same direction as the travelling wave. Fastest type of wave.
S Waves	Travel through solids only. Shakes the Earth vertically (90°angle to the travelling wave). Most damaging type of wave.
Surface waves	They can occur closest to the surface. They travel slower than P and S waves but are more destructive. Love waves: Travel through solids only. Shakes the Earth in the same direction as the travelling wave. Rayleigh waves: Travel through solids and liquids. Shakes the Earth in a rolling motion (like an ocean wave).

Earthquake Secondary Earthquakes

Liquefaction	Solid material changed into a liquid state. Damage to building foundations, results in them sinking.
Landslides and Avalanches	Earthquakes in mountainous regions often cause landslides and avalanches. Steep, unstable slopes are notoriously unstable and vulnerable to landslides.
Tsunamis	Earthquakes occurring underwater can cause the seabed to rise, leading to the displacement of water, producing powerful waves which spread out from the epicentre.

Formation of Tsunamis

1	Large waves caused by the displacement of water triggered by underwater earthquakes, submarine landslides and volcanic eruptions.	
2	In the open ocean, the wave can travel at 500-950km/h and has a wavelength of 200km and a small amplitude (wave height) of 1m.	
3	Closer to land the water gets shallower, causing the waves to increase in size but slow down.	
4	Just before the tsunami reaches the coast, The water withdraws down the shore (drawback).	
5	In Japan 2011, when the tsunami waves reached inland, in some places the waves were 20 metr high. Overall, the tsunami destroyed 200,000 buildings, and killed 19,000 people.	

Volcanic Hazards

Ash cloud	Small pieces of pulverised rock and glass which are thrown into the atmosphere.
Gas	Sulphur dioxide, water vapour and carbon dioxide come out of the volcano.
Lahar	A volcanic mudflow which usually runs down a valley side on the volcano.
Pyroclastic flow	A fast moving current of super-heated gas and ash (1000°C). This travels at 450mph.
Volcanic bomb	A thick (viscous) lava fragment that is ejected from the volcano.

Main Types of Volcanoes

Shield	This type of volcano is almost entirely composed of fluid lava flows . They are found in hot spots or along constructive plate margins . Their eruptions are mostly effusive and predictable .
Composite	Composite volcanoes are created by layers of ash and viscous lava . They can be found along destructive margins and are often steep-sided . They are extremely explosive and unpredictable .

